








I NEED TO STAY HOME IF...

I HAVE A FEVER	I AM VOMITING	I HAVE DIARRHEA	I HAVE A RASH	I HAVE HEAD LICE	I HAVE AN EYE INFECTION	I HAVE BEEN IN THE HOSPITAL
						
Temperature of 100 degrees F or higher	Within the past 24 hours	Within the past 24 hours	Body rash with itching or fever	Itchy head, active head lice and student cannot perform or is distracted from school activities	Redness, itching, and/or pus draining from eye	Hospital stay and/or emergency room visit

I AM READY TO GO BACK TO SCHOOL WHEN I AM...

Fever free without the assistance of medication for 24 hours (i.e., Tylenol, Motrin, Advil)	Free from vomiting for 24 hours and without assistance of anti-vomit meds	Free from diarrhea for 24 hours.	Free from rash, itching or fever. I have been evaluated by my doctor if needed.	Treated with appropriate lice treatment at home.	Free from drainage and/or have been evaluated by my doctor if needed.	Released by my medical provider to return to school.
---	---	----------------------------------	---	--	---	--

If your child has strep throat or another bacterial infection, he/she should stay home until the antibiotic has been given for at least 24 hours and your health care provider has given permission for your child to return to school. We encourage you to seek medical attention when your child is sick and to follow your health care provider's recommendations about returning to school and other activities.

For further information regarding the school exclusion policy please see TXDSHA Communicable Disease Chart at https://dshs.texas.gov/idcu/health/schools_childcar/resources/

NECESITO QUEDARME EN CASA SI...

TENGO FIEBRE	ESTOY VOMITANDO	TENGO DIARREA	TENGO SARPULLIDO	TENGO PIOJOS EN LA CABEZA	TENGO UNA INFECCIÓN DE OJO	ESTUVE EN EL HOSPITAL
						
Temperatura de 100° F o más	Dentro de las últimas 24 horas	Dentro de las últimas 24 horas	Sarpullido corporal con fiebre o picazón	Picazón en la cabeza, piojos activos y el estudiante no puede realizar y se distrae de las actividades escolares	Enrojecimiento, picazón y/o drenaje de pus del ojo	Estancia en hospital y/o visita a sala de urgencias

ESTOY LISTO PARA VOLVER A LA ESCUELA AL SER ...

Libre de fiebre sin la ayuda de medicamentos durante 24 horas (es decir, Tylenol, Motrin, Advil)	Libre de vómitos durante 24 horas. Sin aguda de medicación contra el vómito	Libre de diarrea durante 24 horas.	Libre de sarpullido, picazón o fiebre. He sido evaluado por mi médico si es necesario.	Tratado con tratamiento de piojos apropiado en casa.	Libre de drenaje y/o he sido evaluado por mi medico, si fuera necesario.	Dado de alta por mi proveedor de servicios médicos para regresar a la escuela.
--	---	------------------------------------	--	--	--	--

Si su hijo tiene faringitis estreptocócica u otra infección bacteriana, debe quedarse en casa hasta haberle administrado el antibiótico durante al menos 24 horas y su proveedor de atención médica le haya dado permiso para que regrese a la escuela. Le recomendamos que busque atención médica cuando su hijo esté enfermo y siga las recomendaciones de su proveedor de atención médica sobre regresar a la escuela y otras actividades. Para obtener más información sobre la política de exclusión escolar, consulte el cuadro de enfermedades transmisibles de TXDSHS

https://dshs.texas.gov/idcu/health/schools_childcar/resources/